



# A Perspective on HRT for Women:

Picking Up the Pieces  
After the Women's  
Health Initiative Trial —  
*Part 2*

**George R. Gillson, MD, PhD**  
Rocky Mountain Analytical,  
Calgary, Alberta, Canada

**David T. Zava, PhD**  
ZRT Laboratory,  
Portland, Oregon



*Note: In this article, bioidentical hormone replacement therapy will be referred to as BHRT and conventional hormone replacement therapy will be referred to as CHRT. Progestins will be used throughout this article to refer to synthetic compounds that exert an antiproliferative effect on uterine endometrium. By this definition, progesterone is not a progestin, since it is natural (not synthetic). Testosterone replacement, although relevant, is beyond the scope of our discussion.*

## Progesterone and Progestins

There has been an almost abysmal failure on the part of the medical profession to recognize the difference between progesterone and synthetic progestins. While a detailed comparison of bioidentical progesterone and progestins (synthetic C-19 and C-21 derivatives) will not be provided here, a book by John Lee, MD, a recognized expert on progesterone, discusses this topic in detail.<sup>1</sup> A perusal of the literature indicates that the terms *progesterone* and *progestins* are often used interchangeably. Failure to make the crucial distinction between progesterone and progestins has led us to our present position that compliance with established forms of hormone replacement therapy (HRT) is poor (due to side effects) and that the most commonly prescribed form of combined HRT has been shown to be unsafe over the medium term (5 years) in a large controlled trial.

### Synergy Between Estradiol and Progesterone

One rarely considered aspect of the use of progesterone versus progestins is the synergy between estradiol and progesterone; each is required for the optimal expression of the other's receptors. Progesterone turns on estradiol receptor expression and has also been shown to elevate serum estradiol levels.

Nahoul demonstrated that vaginal administration of progesterone resulted in an elevation of serum estradiol, which was similar in magnitude and duration to the oral administration of 0.5 mg of estradiol.<sup>2</sup> Therefore, concurrent use of progesterone with estrogen (estradiol or estrone) should allow for lower dosing of estrogen. Anecdotally, this has been the experience of many practitioners working with progesterone. In fact, the rule of thumb is to halve the dose of estradiol or estrone when the switch is made from progestins to progesterone. This is Lee's opinion, and he prescribed progesterone skin cream for thousands of women patients for more than 20 years.

### Balance Between Estrone and Estrogen Sulfate Within Tumor Cells

Progesterone also impacts the balance between estrone and estrogen sulfate within tumor cells. It is well known that breast-cancer cells and breast fibroadenomas accumulate large amounts of estrone sulfate, which they then use as a source of estradiol (the sulfate is hydrolyzed to estrone, which is then converted to estradiol).<sup>3</sup> Progesterone, along with some synthetic progestins such as danazol, blocks the conversion of estrone sulfate to estrone and prevents the cells from fueling themselves with estradiol.

### Protection of Uterine Lining

The common feature of progesterone and progestin is their ability to protect the uterine lining from overgrowth/carcinogenesis caused by estrogens. Both progesterone and progestin, when given continuously in opposition to estrogen, produce a dormant or quiescent endometrium. Some critics of progesterone supplementation have focused on studies in which progesterone failed to produce a secretory endometrium. This is not a necessary or even desirable endpoint. A quiescent endometrium is both nonproliferative and nonsecretory. However, an atrophic endometrium (which may arise from progestin therapy) is also undesirable, as it may lead to bleeding and is a degenerate tissue state. Progestins were added to estrogen therapy once it was realized that unopposed estrogen caused uterine cancer, and the focus for many years was only on that aspect. Researchers did not realize that progestins were causing other serious negative effects, such as increased risk of breast cancer and cardiovascular disease.

### Impact on Cardiovascular Function

Many human and animal studies have indicated that estradiol does exert positive effects on parameters influencing cardiac disease.<sup>4-6</sup> These effects include vasodilation,<sup>4</sup> reduction of vascular proliferative and inflammatory responses,<sup>5</sup> and decreased sympathetic activity.<sup>6</sup>

However, the Women's Health Initiative (WHI) trial indicated that oral conjugated estrogens and oral medroxyprogesterone acetate increased cardiac risk. This apparent paradox is resolved by realizing that there are many studies that indicate that medroxyprogesterone acetate is detrimental to cardiovascular function. Clarkson has reviewed the adverse effects of medroxyprogesterone acetate on cardiovascular health in some detail.<sup>7</sup> For examples of such studies, see sidebar.

### Studies Showing Adverse Effects of Medroxyprogesterone Acetate on Cardiovascular Health

- A study by Miyagawa demonstrated that estradiol plus medroxyprogesterone failed to prevent coronary vasospasm in response to serotonin plus a thromboxane A2 mimetic in rhesus monkeys, whereas estradiol plus progesterone was protective against vasospasm.<sup>8</sup>
- A study conducted by Wakatsuki demonstrated that medroxyprogesterone acetate dose-dependently inhibited improvement in flow-mediated brachial artery vasodilation seen in postmenopausal women who were given oral estrogen.<sup>9</sup>
- Kojima demonstrated that medroxyprogesterone acetate interferes with the assemblage of cholesterol and phospholipids into high-density lipoprotein (HDL) particles by apolipoprotein A-1.<sup>10</sup> Although progesterone also decreased HDL, the effect was smaller.
- In the Postmenopausal Estrogen/Progestin Interventions (PEPI) trial, it was found that at the end of 3 years, oral estrogen increased HDL cholesterol (net increase 7%), but this increase was considerably reduced (net increase 2%) when medroxyprogesterone acetate was added.<sup>11</sup>

On the other hand, evidence that progesterone exerts beneficial effects on cardiovascular function is accumulating. For examples of such studies, see sidebar.

### Studies Showing Progesterone's Beneficial Effects on Cardiovascular Function

- With monkeys as test subjects, Adams demonstrated that, whereas medroxyprogesterone acetate substantially blunted the reduction in plaque induced by conjugated estrogens, progesterone had no effect on the reduction in atherosclerotic plaque induced by estradiol.<sup>12,13</sup>
- A recent study by Rosano has shown that vaginally administered progesterone had a beneficial effect on cardiovascular health. The time to ischemia on a treadmill was increased in postmenopausal women who had cardiovascular disease.<sup>14</sup>
- In another recent study, Molinari indicated that an intravenous infusion of progesterone increased blood flow in porcine mesenteric, iliac and renal arteries via the release of nitrous oxide.<sup>15</sup>
- With the use of cynomolgus monkeys, a study has shown that medroxyprogesterone acetate blunts the estradiol-induced release of nitrous oxide, a potent vasodilator.<sup>16</sup>

Although it is reasonably clear that some progestins may be deleterious to cardiac health, a recent review article points out that not all progestins exert equivalent effects on HDL, triglycerides, and other relevant parameters.<sup>17</sup> In fact, a recent small study by Sander-son demonstrated that estradiol plus

norethindrone acetate improved exercise tolerance, increased time to ST segment elevation on a treadmill and reduced the total number of ischemic events during 24-hour ambulatory electrocardiographic monitoring in patients with angina.<sup>18</sup> Pharmaceutical companies will eventually find progestins that are safer and that

will more closely mimic progesterone. However, if we do find these synthetic molecules, they will likely be so close in effect to progesterone that we might as well use progesterone.

## Bioidentical Progesterones

### Oral Bioidentical Progesterone

Bioidentical progesterone in the form of off-the-shelf progesterone capsules (in peanut oil), as well as compounded micronized progesterone capsules, has already been accepted by many physicians; and various publications support the ability of oral progesterone to produce a quiescent endometrium. For examples of studies investigating bio-identical progesterone, see sidebar.

### Studies of Bioidentical Progesterone

- In a 12-month study of 10 women, Hargrove showed that oral micronized progesterone (100 mg/twice a day or 100 mg/mornings and 200 mg/evenings) in conjunction with oral micronized estradiol (0.35 mg/twice a day or 0.35 mg/mornings and 0.7 mg/evenings) resulted in a quiescent endometrium.<sup>19</sup> At 6 months, there was no uterine bleeding in this group, whereas four of five women in the same study who were on 0.625 mg/day conjugated estrogens plus medroxyprogesterone acetate (10 mg/day for 10 days each month) continued to have withdrawal bleeding throughout the study.
- In a study by Gillet,<sup>20</sup> 98 women received transdermal estradiol (1.5 mg/day) plus 100 mg/day oral micronized progesterone (21 to 25 days/month). Endometrial biopsies were performed after a minimum of 6 months of therapy. No endometrial hyperplasia was found in any woman after a minimum of 6 months of supplementation, and 92% of women were amenorrheic after 6 months.<sup>21</sup> No endometrial biopsy could be classified as proliferative, and the mean number of mitoses/1000 cells was 0.5 (very low).
- In the much longer PEPI trial, oral micronized progesterone (200 mg/day) was as effective as Provera (10 mg/day) in suppression of endometrial hyperplasia, when used in conjunction with 0.625 mg/day of conjugated equine estrogen.<sup>11</sup>

# HT

*doesn't have to be  
confusing...*

### Eliminate Guesswork with ZRT Lab's Hormone Test Kit

*Saliva and bloodspot hormone testing accurately detects hidden hormone imbalances and helps physicians prescribe safe, individualized hormone supplementation.*


#### Simple and Private

- Provider and patient determine hormones to test based on symptom checklist.
- Provider stocks and dispenses home collection kit indicating hormones to be tested.
- Patient collects specimen at home in tube and mails sample and documentation to ZRT Lab.
- ZRT analyzes hormone specimen, relating symptoms and hormone usage to patient test results for a more comprehensive evaluation.
- Patient, physician and pharmacist determine safe individualized HT to restore hormone balance and relieve symptoms.

**NEW** - Blood Spot Male Hormone Profile (I & II) Test for PSA, Testosterone, SHBG and IGF-1

## 866-600-1636

www.salivatest.com www.bloodspottest.com  
1815 NW 169th Place, Suite 5050 • Beaverton, OR 97006



Track patient progress  
with comparative  
history reports

Monitor and adjust  
hormone levels  
accordingly

ZRT  
LABORATORY

Despite its efficacy, oral administration of bioidentical progesterone has several potential drawbacks. Since progesterone is suspended in peanut oil, it poses a problem for individuals who are allergic to peanuts. Oral progesterone is not well tolerated by some women, with side effects that include nausea, breast swelling, dizziness, drowsiness, and depression. As is the case for oral estradiol, progesterone undergoes a substantial amount of first-pass gut and liver metabolism.<sup>2,21</sup> It is progesterone metabolites such as allopregnanolone and deoxycorticosterone that are thought to be responsible for the side effects.<sup>22-24</sup>

### Topical Bioidentical Progesterone

Progesterone skin cream has been widely used over the past 20 years as an alternative to oral progesterone and

progestins. Leonetti reported that progesterone cream (20 mg/day) given to postmenopausal women for 1 year (without estrogen replacement therapy) relieved vasomotor symptoms in 25 of 30 women (83%), versus 5 of 26 women (19%) given placebo.<sup>25</sup> Progesterone is necessary to prime estradiol receptors and may elevate endogenous estradiol levels; therefore, provision of progesterone alone is able to affect symptoms such as vasomotor complaints thought to be related primarily to low estrogen levels.


Many physicians very rightly question whether enough progesterone can be absorbed through the skin to protect the endometrium. There are also concerns about the standardization of over-the-counter progesterone creams, although this is not an issue for compounded products. Considering the findings of

the following studies can help in drawing conclusions about these issues.

Levine measured total serum progesterone levels (via liquid chromatography/mass spectroscopy) after ingestion of 100 mg of progesterone in peanut oil. A peak serum level of 2.4 ng/mL was achieved 1 hour after ingestion, followed by a rapid decline to baseline.<sup>21</sup> However, the previously mentioned study by Gillet showed that 100 mg/day oral progesterone is sufficient to produce a histologically quiescent endometrium.<sup>20</sup> Therefore, we can conclude that an average serum progesterone level of somewhat less than 2.4 ng/mL is likely sufficient to stabilize the endometrium, depending on the dose of estrogen being administered. Luteal-phase progesterone levels (10 to 30 ng/mL) may be required to produce luteal phase histology (ie, a secretory

## AirClean Systems

## PowderSafe™



### Controlling Occupational Exposure to Hazardous Powder and Particulate

#### Features:

- No Ductwork Required - Portable
- Constant monitoring and automatic adjustment of airflow
- Continuous monitoring of HEPA filter loading
- Available in 32-inch and 48-inch widths
- Real Time Airflow Display
- Factory leak tested and certified

Independently certified to current compounding procedures.

Website: [www.aircleansystems.com](http://www.aircleansystems.com)

Toll Free: (800) 849-0472

E-mail: [info@aircleansystems.com](mailto:info@aircleansystems.com)

endometrium); but, evidently, daily exposure to a much lower progesterone level can stabilize the endometrium (depending on the estrogen drive that has to be overcome).

We must also consider whether transdermal delivery of progesterone can produce serum total progesterone levels in this range (1 to 3 ng/mL). Transdermal delivery includes intranasal and vaginal sites, as well as peripheral skin delivery. In a study by Cicinelli, the mean serum progesterone level in eight women after 9 days of nasally administered progesterone (11 mg/three times a day) was 3.9 ng/mL (a sixfold increase from baseline).<sup>26</sup> A mean serum progesterone level of 2.4 ng/mL was seen after 13 days of administration of vaginal progesterone gel (45 mg/day) to 14 women.<sup>27</sup> In both these studies, estrogens had been administered for 28 days prior to starting progesterone; and, in both these studies, secretory endometrium was found on biopsy. Therefore, these levels are actually likely higher than that needed to simply stabilize the endometrium in a quiescent mode.

Several studies also show the results of serum progesterone levels achieved after application of progesterone to the skin (see sidebar).

## Bio-Identical HRT Quick Reference CD from IJPC

- Permanent storage and easy retrieval of articles and formulations.
- Print articles in color from the pdf file for your marketing needs.
- Eliminate long searches through past issues for articles/formulations.

**Only \$65** (plus \$2.50 for S&H).

### More than 30 feature articles including:

- Natural Hormone Replacement Therapy: What It Is and What Consumers Really Want
- The Science Behind Bioidentical Hormone Replacement Therapy
- "Natural" Isomolecular Hormone Replacement: An Evidence-Based Medicine Approach
- Essential Truths About HRT
- Using Serum Hormone Analysis to Develop Hormone Replacement Therapy Regimens
- Finding the Balance
- Natural HRT: Corrective Measures When the System Goes Down
- Estrogen and Breast Cancer: Is There a Link?
- Estrogen in the Treatment and Prevention of Alzheimer's Disease
- Bioidentical Topical Progesterone: A Survey of Patients
- The Role of Natural Progesterone in Natural Hormone Replacement Therapy
- Treating Andropause: Prohormones and Hormone Metabolic Modifiers
- Compounded Testosterone Gels: A Guide for Clinicians and Pharmacists
- Testosterone Deficiency in Men: New Treatments for Andropause
- Interview on HRT: Christiane Northrup
- Interview: Katharina Dalton: Progesterone and Related Topics
- Long-Term Effects of Topical Progesterone Cream Applications
- Testosterone Supplementation in a Vasectomy Patient
- Planning and Marketing a Natural HRT Patient Seminar
- HRT and the Continuing Education of Health Care Professionals
- Bioidentical Hormones in the Treatment of Osteoporosis
- Infertility Care and the Role of the Compounding Pharmacist

### And 50 formulations!

For a complete list of contents including formulations, visit our website.

**To order, call 888-588-4572 or visit our website at [www.ijpc.com/products](http://www.ijpc.com/products).**

### Studies Showing Results With Topical Progesterone

- Mircioiu compared serum progesterone time profiles after a one-time application of 90 mg progesterone in gel (n=9), emulsion (n=16), and lipophilic base cream (n=7).<sup>28</sup> In all cases, serum levels peaked 3 to 4 hours after application and declined gradually over the ensuing 72 hours. Peak levels were in the range of 1 to 2.2 ng/mL.
- Burry administered progesterone cream to six women for 28 days (14 days:30 mg/day, 14 days:30 mg/twice a day) and measured serum progesterone time profiles (eight points in 24 hours) on each of 5 days (days 1, 8, 15, 22, and 29).<sup>29</sup> After 1 week of use, five of the six women sustained a progesterone level of at least 1 ng/mL for at least 8 hours after application. On day 29, the average progesterone level was 2.2 ng/mL. Serum progesterone level was dose related and consistent for each woman, although interindividual results varied by a factor of two.
- Carey measured serum progesterone time profiles after 6 weeks of progesterone cream (20 mg/twice a day, n=10; and 40 mg/day, n=10).<sup>30</sup> Both regimens resulted in average progesterone levels of roughly 0.7 ng/mL over 24 hours. Peak levels were approximately 1 ng/mL.
- O'Leary measured serum progesterone levels at several time points after a one-time application of progesterone cream (64 mg) in six premenopausal and six postmenopausal women.<sup>31</sup> In the premenopausal group, serum progesterone did not increase from an average baseline of 6.6 ng/mL. Mean serum progesterone 4 hours after application increased slightly to 1.1 ng/mL from 0.75 ng/mL at baseline in the postmenopausal women.
- Wren administered cyclic (14 days/month) progesterone skin cream to postmenopausal women in varying doses for 3 months.<sup>32</sup> Nine women received 16 mg/day, eight received 32 mg/day, and ten received 64 mg/day. Curiously, the average serum progesterone level achieved was only 0.4 ng/mL, and one has to question the degree of absorption of the cream used in that study.

The majority of published literature indicates that transdermally delivered progesterone can produce serum total progesterone levels in the range of 1 to 3 ng/mL, which are likely to produce a quiescent endometrium. A definitive study combining progesterone cream administration, measurement of serum total progesterone levels, and endometrial biopsy has not been done. However, a summary of a 4-week trial of progesterone cream with serial endometrial biopsies on 58 women pretreated with conjugated estrogens has been published.<sup>33</sup> Twice-daily vaginal and topical progesterone creams of varying strengths were compared with placebo. All routes and concentrations of progesterone supplementation produced a significant decrease in the degree of endometrial proliferation induced by conjugated estrogens (0.625 mg/day) compared with placebo.

There are several additional lines of evidence that indicate that transdermal progesterone cream is delivered to tissue in

appreciable amounts. The first body of evidence comes from salivary hormone measurements. When a steroid hormone skin cream, such as progesterone, is rubbed on a relatively small area of skin, the salivary progesterone level increases dramatically within several hours and often by several orders of magnitude. For a given individual, this elevation is reproducible and dose dependent, although the degree of elevation varies between individuals for a given dose and site of application. A full explanation of the mechanism by which steroids enter saliva under these circumstances is beyond the scope of this article. Suffice it to say that transdermal application of steroids likely enriches the concentration of steroids in red blood cell (RBC) membranes. Due to the tight fit as the RBCs wedge their way through the capillaries, steroid hormones may be efficiently transferred directly to tissue by partitioning into the membranes of the capillary endothelial cells. Therefore, a salivary hormone level reflects delivery of hormone to cell membranes (tissue), unlike a blood level, which measures the amount of hormone that has not been delivered to tissue.

Chang examined the effect of topical progesterone and estradiol on breast tissue uptake and cell proliferation in women scheduled for biopsies or reduction mammoplasties.<sup>34</sup> Women were treated for 10 to 13 days with a placebo gel or gels that contained 1 mg estradiol, 25 mg progesterone, or a combination of estradiol and progesterone. Biopsy samples were taken, and half were analyzed for estradiol and progesterone content; the other half were analyzed by a pathologist for breast-cell proliferation. Estradiol and progesterone increased 200-fold and 100-fold with respective topical delivery, demonstrating tissue uptake. Estradiol increased breast-cell proliferation twofold, and progesterone decreased breast-cell proliferation from baseline but also suppressed proliferation activated by estradiol. Serum estradiol or progesterone did not increase significantly in any of the treatments, despite a remarkable tissue uptake and biological response. These results support the notion that tissue uptake and response can occur with topical hormone delivery without a noticeable effect on serum levels. A 25-mg dose of progesterone dramatically increases salivary progesterone to levels of 0.5 to 3 ng/mL at 12 to 24 hours after supplementation and peak levels (3 hours after application) above 20 ng/mL. Therefore, it is reasonable to conclude that salivary measurements are more meaningful than blood measurements in the context of transdermal hormone supplementation, since the former more closely reflects hormone delivery to tissue.

Optimum results with progesterone skin cream will likely be achieved with standardized products delivered in reproducible fashion with metered-dose delivery. Women who have not undergone a hysterectomy and use progesterone and estrogen(s) should be monitored with ultrasound and/or endometrial biopsy as indicated to evaluate the status of the endometrium. Practitioners cannot assume that, because progesterone is more natural, it is automatically protecting the endometrium over the long term.

Oral progestins exert a much stronger suppressant effect on the endometrium, which reduces our concerns about the endometrium. This is achieved at the expense of side effects; increased risk of cancer and cardiovascular and thromboembolic disease; and induction of an atrophic, degenerative tissue state, with increased risk of breakthrough bleeding. The published literature indicates that we cannot assume anything about the protection of the endometrium with transdermally administered progesterone, but we can have the expectation that it will do the job, although thought and attention are required. Certainly larger, more definitive trials of progesterone skin cream are needed; and at least one large, long-term study that compares progesterone skin cream/oral estrogen to oral progestin/oral estrogen is in progress. A protocol for a reasonable approach to combined HRT that employs progesterone skin cream, based on the foregoing discussion, has been included (see sidebar, p. 336).

### Conclusion

We have spent 20 years researching both estrogen only and estrogen/progestin HRT. Estrogen replacement has been shown to relieve menopausal symptoms, improve bone and skin health and diminish long-term morbidity from

## Kalchem International

Chemicals & Compounding Supplies

- Family owned and operated with over 50 years of combined experience in the pharmaceutical/healthcare industry.
- We import a wide range of products from around the world.
- Competitive pricing.
- Kalchem can source anything from chemicals and capsules to equipment. Schedule II and III narcotics also available.
- All chemicals quarantined and tested for correctness and potency.
- All chemicals meet or exceed USP standards.

**We are a full line chemical company that can provide you with all your pharmaceutical compounding needs.**

**Call Toll Free: 888-298-9905, or  
Order from our secure website:  
[www.kalcheminternational.com](http://www.kalcheminternational.com)**

224 South Main Street, Lindsey, OK 73052

### Protocol for Combined HRT Using Progesterone Skin Cream

1. Compounded progesterone cream, 20 to 30 mg/twice daily or 25 days/month, applied anywhere but the abdomen (to avoid first-pass liver metabolism), with metered-dose delivery (eg, syringe or jar with détente delivery system);
2. Pre- and postapplication serum/salivary progesterone levels (post: 8 hours after cream application), after at least one cycle (25 to 30 days) of progesterone use, aiming for 1 to 3 ng/mL serum and 1 to 5 ng/mL salivary progesterone;
3. Estrogen replacement therapy 25 days/month concurrent with progesterone cream at a dose that does not result in withdrawal bleeding or spotting, such as:  
0.5 to 1.0 mg estriol (oral or cream) twice daily, plus 0.05 to 0.1 mg estradiol (oral or cream) twice daily;  
or 0.075 to 0.15 mg estradiol cream twice daily;  
or estradiol slow-release patch 0.025 to 0.1 mg/24 hours;
4. Transvaginal ultrasound before institution of therapy and at 6-month intervals for the first 12 to 18 months, with endometrial biopsy as indicated for bleeding or spotting persisting after 3 months of initiation of therapy; and
5. Periodic monitoring of bone density, with adjustment of therapy as indicated.

cardiovascular disease. However, these benefits are accompanied by increased risks of breast cancer, venous thromboembolic disease, gallbladder disease, and endometrial hyperplasia/cancer. Addition of a progestin mitigates against endometrial problems but increases other risks, as demonstrated conclusively by the results of the WHI trial and other studies. Side effects from progestins seriously compromise compliance.

Patients have become increasingly aware of hormone replacement with bioidentical hormones and are coming to their physicians on a daily basis with questions about this more natural HRT option. We shouldn't hedge and turn our backs on patients suffering from hormonal deficiencies while we wait for large trials to be conducted. Moreover, we are not sure if large trials will be conducted or who will fund them, since bioidentical hormones are not patentable. We should go back to first principles, think for ourselves, incorporate what we have learned from more than 30 years of steroid hormone research, and make our best call. We have good indications from the literature about what types of estrogens we should use (estradiol and estriol), and we realize that transdermal delivery of hormones offers advantages over oral delivery. If oral delivery is used, we need to rethink dosing and be aware of the impact on other relevant parameters such as hormone-binding

## COMPOUND WITH CONFIDENCE...

The products you compound enhance the quality of life for patients but can endanger your employees' health and your pharmacy.

MIC COMPOUNDING SYSTEMS provide safe containment levels acceptable for compounding hormones and other potent products. Our SYSTEMS assure a regulatory/acceptable environment that meets the requirements of OSHA, FDA and state agencies for parenteral and solid dosage products. Our ergonomically integrated systems include both equipment and enclosures.


MIC  
COMPOUNDING  
SYSTEMS

DON'T TAKE CHANCES WITH YOUR FUTURE.



**CONTAINMENT TECHNOLOGIES GROUP, INC.**  
110329 Vandergriff Road ■ Indianapolis, IN 46239  
(317) 862-5945 ■ [mic@mic4.com](mailto:mic@mic4.com) ■ [www.mic4.com](http://www.mic4.com)

Please visit  
[www.mic4.com](http://www.mic4.com) for  
recent test results



proteins and other hepatic proteins. There is evidence that we should avoid certain progestins, and there is strong evidence to support both the oral and transdermal use of bioidentical progesterone. Through testing modalities such as saliva, we are able to more closely monitor the delivery of hormone to tissue.

With good follow-up, good clinical judgment and good laboratory back-up, we have enough information about hormone replacement to offer BHRT as a viable option. It might take a little more thought on our part, but at least we will not be leaving the decision about HRT up to our patients (we know that many of them already vote with their feet). As time goes on, we need to analyze new information and modify our approach as indicated.

## References

- Lee J, Hanley J, Hopkins V. *What Your Doctor May Not Tell You About Premenopause: Balance Your Hormones and Your Life From Thirty to Fifty*. New York:Warner Books, Inc.; 1999.
- Nahoul K, Dehennin L, Jondet M et al. The metabolism of estradiol: Oral compared to intravenous administration. *J Steroid Biochem* 1985;23:1065-1070.
- Pasqualini JR, Chetrite G, Nestour EL. Control and expression of oestron sulphatase activities in human breast cancer. *J Endocrinol* 1996;150 Suppl:S99-105.
- Komesaroff PA, Black CV, Westerman RA. A novel, nongenomic action of estrogen on the cardiovascular system. *J Clin Endocrinol Metab* 1998;83:2313-2316.
- Dubey RK, Jackson EK, Gillespie DG et al. Clinically used estrogens differentially inhibit human aortic smooth muscle cell growth and mitogen-activated protein kinase activity. *Arterioscler Thromb Vasc Biol* 2000;20:964-972.
- Vongpatanasin W, Tuncel M, Mansour et al. Transdermal estrogen replacement therapy decreases sympathetic activity in postmenopausal women. *Circulation* 2001;103:2903-2908.
- Clarkson TB. Progestogens and cardiovascular disease. A critical review. *J Reprod Med* 1999;44(2 Suppl):180-184.
- Miyagawa K, Rosch J, Stanczyk F et al. Medroxyprogesterone interferes with ovarian steroid protection against coronary vasospasm. *Nat Med* 1997;3:324-327.
- Wakatsuki A, Okatani Y, Ikenoue N et al. Effect of medroxyprogesterone acetate on endothelium-dependent vasodilation in postmenopausal women receiving estrogen. *Circulation* 2001;104:1773-1778.
- Kojima K, Abe-Dohmae S, Arakawa R et al. Progesterone inhibits apolipoprotein-mediated cellular lipid release: A putative mechanism for the decrease of high-density lipoprotein. *Biochim Biophys Acta* 2001;1532:173-184.
- The Writing Group for the Postmenopausal Estrogen/Progestin Interventions (PEPI) Trial. Effects of estrogen or estrogen/progestin regimens on heart disease risk factors in postmenopausal women. *JAMA* 1995;273:199-208.
- Adams MR, Kaplan JR, Manuck SB et al. Inhibition of coronary artery atherosclerosis by 17-beta estradiol in ovariectomized monkeys: Lack of an effect of added progesterone. *Arteriosclerosis* 1990;10:1051-1057.
- Adams M, Register T, Golden D et al. Medroxyprogesterone acetate antagonizes inhibitory effects of conjugated equine estrogens on coronary artery atherosclerosis. *Arterioscler Thromb Vasc Biol* 1997;17:217-221.
- Rosano GM, Webb CM, Chierchia S et al. Natural progesterone, but not medroxyprogesterone acetate, enhances the beneficial effect of estrogen on exercise-induced myocardial ischemia in postmenopausal women. *J Am Coll Cardiol* 2000;38:2154-2159.
- Molinari C, Battaglia A, Grossini E et al. Effect of progesterone on peripheral blood flow in prepubertal female anesthetized pigs. *J Vasc Res* 2001;38:569-577.
- Williams JK, Honore EK, Washburn SA et al. Effects of hormone replacement therapy on reactivity of atherosclerotic coronary arteries in cynomolgus monkeys. *J Am Coll Cardiol* 1994;24:1757-1761.
- Godsland IF. Effects of postmenopausal hormone replacement therapy on lipid, lipoprotein, and apolipoprotein (a) concentrations: Analysis of studies published from 1974-2000. *Fertil Steril* 2001;75:898-915.
- Sanderson JE, Haines CJ, Yeung L et al. Anti-ischemic action of estrogen-progestogen continuous combined hormone replacement therapy in postmenopausal women with established angina pectoris: A randomized, placebo-controlled, double-blind, parallel-group trial. *J Cardiovasc Pharmacol* 2001;38:372-383.
- Hargrove JT, Eisenberg E. Menopause. *Med Clin North Am* 1995;79:1337-1356.
- Gillet JY, Andre G, Faguer B et al. Induction of amenorrhoea during hormone replacement therapy: Optimal micronized progesterone dose. A multicenter study. *Maturitas* 1994;19:103-115.
- Levine H, Watson N. Comparison of the pharmacokinetics of crinone 8% administered vaginally versus Prometrium administered orally in postmenopausal women. *Fertil Steril* 2000;73:516-521.

## Is your compounding base free?



...free of  
dyes, perfume, fragrance  
and masking fragrance,  
lanolin, parabens and  
formaldehyde

- non-irritating
- non-greasy
- non-comedogenic
- quickly absorbed
- easy to apply
- widely compatible
- recommended by dermatologists
- stable
- oil-in-water emulsion
- produces elegant products
- easy equipment cleanup
- cosmetically acceptable
- competitively priced

To receive a free lb. jar  
or for more information, call 1-800-325-8232

PHARMACEUTICAL SPECIALTIES, INC.  
ROCHESTER, MN 55903-0298



22. Arafat ES, Hargrove JT, Maxson WW et al. Sedative and hypnotic effects of oral administration of micronized progesterone may be mediated through its metabolites. *Am J Obstet Gynecol* 1988;159:1203-1209.
23. Freeman EW, Purdy RH, Coutifaris C et al. Anxiolytic metabolites of progesterone: Correlation with mood and performance measures following oral progesterone administration to healthy female volunteers. *Neuroendocrinology* 1993;58:478-484.
24. Ottoson UB, Carlstrom K, Damber JE et al. Serum levels of progesterone and some of its metabolites including deoxycorticosterone after oral and parenteral administration. *Br J Obstet Gynaecol* 1984;91:1111-1119.
25. Leonetti HB, Longo S, Anasti JN. Transdermal progesterone cream for vasomotor symptoms and postmenopausal bone loss. *Obstet Gynecol* 1999;94:225-228.
26. Cicinelli E, Cignarelli M, Resta L et al. Effects of the repetitive administration of progesterone by nasal spray in postmenopausal women. *Fertil Steril* 1993;60:1020-1024.
27. Fanchin R, De Ziegler D, Bergeron C et al. Transvaginal administration of progesterone. *Obstet Gynecol* 1997;90:396-401.
28. Mircioiu C, Perju A, Griu E et al. Pharmacokinetics of progesterone in postmenopausal women: 2. Pharmacokinetics following percutaneous administration. *Eur J Drug Metab Pharmacokinet* 1998;23:397-402.
29. Burry KA, Patton PE, Hermsmeyer K. Percutaneous absorption of progesterone in postmenopausal women treated with transdermal estrogen. *Am J Obstet Gynecol* 1999;180:1504-1511.
30. Carey BJ, Carey AH, Patel S et al. A study to evaluate serum and urinary hormone levels following short and long term administration of two regimens of progesterone cream in postmenopausal women. *BJOG* 2000;107:722-726.
31. O'Leary P, Feddema P, Chan K et al. Salivary, but not serum or urinary levels of progesterone are elevated after topical application of progesterone cream to pre- and postmenopausal women. *Clin Endocrinol (Oxf)* 2000;53:615-620.
32. Wren BG, McFarland K, Edwards L et al. Effect of sequential transdermal progesterone cream on endometrium, bleeding pattern, and plasma progesterone and salivary progesterone levels in postmenopausal women. *Climacteric* 2000;3:155-160.
33. Anasti JN, Leonetti HB, Wilson KJ. Topical progesterone cream has antiproliferative effect on estrogen-stimulated endometrium. *Obstet Gynecol* 2001;97(Suppl 1):S10.
34. Chang KJ, Lee TT, Linares-Cruz G et al. Influences of percutaneous administration of estradiol and progesterone on human breast epithelial cell cycle in vivo. *Fertil Steril* 1995;63:785-791.

Address correspondence to: George Gillson, MD, PhD, Rocky Mountain Analytical, Unit A, 253147 Bears paw Road, NW, Calgary, Alberta, T3L 2P5, Canada. E-mail: [gillson@rmlab.com](mailto:gillson@rmlab.com) ■


## The Paddock Solid-to-Liquid Solution

The physical and chemical properties of Ora-Plus<sup>®</sup> and Ora-Sweet<sup>®</sup> help ensure the stability of compounded oral liquids.

Over 40 formulations have been independently tested.

Call 1-800-328-5113 to receive copies of the studies or visit our website, [www.paddocklabs.com](http://www.paddocklabs.com).

### PADDOCK COMPOUNDING



SMART ALTERNATIVES

**Paddock**  
Laboratories, Inc.

## PATIENT HANDOUT

# Hormone Replacement Therapy

### Recent Studies

Many women and their physicians have been shocked to learn that conventional hormone replacement therapy (HRT) can actually increase the chance of getting a major disease. The most recent controversial Women's Health Initiative trial was abruptly halted in July 2002 because it showed that conventional HRT, using synthetic hormones such as Premarin, Prempro, Premphase, Provera, Cycrin, Amen, or any medication containing conjugated equine estrogens (made from horse urine) and a synthetic (manmade) progestin, medroxyprogesterone acetate, increased the chance of some major diseases, as follows:

- A 41% increase in strokes,
- A 29% increase in heart attacks,
- A 26% increase in breast cancer,
- A 22% increase in total cardiovascular disease,
- A doubling of the rate of blood clots, and
- A possible contribution to Alzheimer's disease.



### Options

Many women taking conventional HRT feel they are faced with only two options: continue to take the synthetic hormones and hope for the best, or stop taking hormones and possibly experience unpleasant symptoms such as hot flashes, vaginal dryness, bloating, forgetfulness, mood swings, night sweats, irritability, breast soreness and weight gain, to name a few.

A third choice you may not be aware of is bioidentical HRT. This therapy differs from conventional HRT in several ways.

#### Bioidentical HRT

1. Uses hormones with the same structure as the hormones produced by the human body (synthetic hormones have the structure appropriate for horses, with only approximately 30% identical to human hormones);
2. Is prepared by a compounding pharmacist and not available from a drug manufacturer;
3. Uses progesterone, not progestin, and different estrogens (estriol, estradiol and estrone) either as a single-ingredient or multiple-ingredient prescription;
4. Is individualized or customized to each woman's needs, symptoms, hormone levels, etc., and can be easily adjusted; and
5. Is prepared in a dosage form, ie, troche, capsule, cream, or suppository, based on what your doctor recommends and what you prefer.

---

**W**hether you're looking for relief from symptoms caused by hormone imbalance for the first time or trying to decide if you should continue to take the hormones you've been taking, you should be aware of all the options. Your compounding pharmacist can provide more information about the option of bioidentical HRT. Working with you and your physician, he or she can determine what bioidentical HRT is right for you.

If you are interested in learning more about bioidentical HRT, please contact a compounding pharmacist for further information. It is possible you can work with him or her and your physician to identify a therapy that is just right for you.